



acadian



NUTRITIONAL PRODUCTS FOR QUALITY CROPS[®]

ROOT AND SHOOT BUILDER

0.1 | 0.5 | 1.5

GUARANTEED MINIMUM ANALYSIS

NITROGEN (N) AS ORGANIC	0.1%
PHOSPHORUS (P) AS WATER SOLUBLE	0.4%
POTASSIUM (K) AS OXIDES	1.2%
POTASSIUM (K) AS ORGANIC	0.1%
TOTAL POTASSIUM	1.3%
DERIVED FROM ASCOPHYLLUM NODOSUM	

Manufactured by:



Acadian AgriTech
30 Brown Avenue, Dartmouth
Nova Scotia, Canada, B3B 1X8
Tel.: +1-902-468-2840
Fax: +1-902-468-3474
www.acadianagritech.com

Distributed by:



1 800 634 204
info@ocp.com.au

NET CONTENT:
10 Litres

PRODUCT OF CANADA 

A1244a.0811



DIRECTIONS FOR USE

Mixing Instructions: Fill half of the mixing container with water and gradually add ACADIAN[®] with constant and thorough mixing. Bring to final volume with continued mixing.

Foliar Applications: Fill half the spray tank with water and begin agitating while gradually adding the recommended volume of ACADIAN[®] to the remainder of the water and spray solution. Higher dilution rates are recommended for the best crop coverage. Product should be applied in 500 to 2,000 litres of water per hectare for ground sprayers or 100 to 200 litres of water per hectare for low volume ground sprayers or aerial applications.

Soil Applications: Apply ACADIAN[®] in furrow-run irrigation water by adding solution into the irrigation water. Each nozzle should be calibrated to drip at a constant rate into the irrigation water. For best results, apply through an irrigation system.

Application Rates: Suggested rates and dosages are approximate and may vary depending on crop, climatic region, soil type and fertility. It is preferable to increase the frequency of the applications rather than the concentration of the solution.

Fruit Production: Apply ACADIAN[®] at 2 to 4 litres per hectare per application at the following growth stages:

- 1) Early spring growth
- 2) Pre bloom
- 3) Post bloom
- 4) Early sizing stages
- 5) Late sizing stages

Vegetable Production: Apply ACADIAN[®] at 2 to 4 litres per hectare per application at the following growth stages:

- 1) Transplant drench
- 2) 10 to 14 days post transplant
- 3) 3 to 4 weeks intervals during growing season (Pre/post bloom)
- 4) Fruit/root/tuber development

Ornamentals, Flowers & Gardens: Apply ACADIAN[®] at a rate of 50 millilitres per 100 square meters in a 1:500 dilution.

General Home Garden Use: Make a 0.5% to 2% solution of ACADIAN[®], apply liberally every 10 to 14 days.

GENERAL INFORMATION

ACADIAN[®] is water soluble and is suitable for use in a conventional foliar and irrigation water applications system.

SHAKE BEFORE USE

COMPATIBILITY

ACADIAN[®] is compatible with most insecticides, fungicides and fertilizers. When interaction is unknown, a "jar" compatibility test is recommended.

STORAGE

Store in cool dry place away from direct sunlight.

CAUTION

Keep out of reach of children.

Note: Foliar treatments of ACADIAN[®] may supply only a portion of the major and minor nutrients (nitrogen, phosphorous and potassium) required for successful crop production. This product is recommended as a supplementary plant nutrient source to be used as part of a comprehensive fertilizer program based on soil fertility and plant nutritional demands.

Contact your local dealer or advisor for specific use recommendations. Application rates may vary based on region, soil conditions, economics, climate, etc.

DISTRIBUTED BY:



1 800 634 204
info@ocp.com.au



8 72488 00130 1

A1244b.0811