




## Seasol Plus Calcium

**Seasol Plus Calcium** is a calcium fortified Seasol concentrate. This product has all the benefits of Seasol, with calcium for improving cell-wall integrity in developing fruit and leaves. Improved fruit firmness and colour at harvest. Better quality retention through packing and cool-storage.

PACK SIZES



**INSTRUCTIONS FOR USE:** 1. **Seasol Plus Calcium** must be diluted in water. Use sufficient water to achieve adequate coverage when applying as a foliar spray. 2. Shake or agitate before use. Maintain agitation during dilution and application if possible. 3. Climatic stress - use 5/10L / ha 3-7 days prior to and after extreme stress events (e.g. heatwave and frost) to mitigate crop damage and assist recovery.

Category	Crop	Rate	Application / Critical Comments
<b>Nut Trees</b>	Almonds, Hazelnuts, Macadamias, Pecans	5-10L/ha or 1ml per sq mtr of plant area	Planting - drench or water in with 1:400 dilution. Apply from green tip/first growth flush. Recommend 5 applications at critical growth stages inc post harvest. Use higher rate for larger trees
<b>Deciduous Fruit Trees</b>	Pome Fruits, Stone Fruits and other deciduous species	5-10L/ha or 1ml per sq mtr of plant area	Planting - drench or water in with 1:400 dilution. Apply at each growth flush or 5 applications at critical growth stages and post harvest. Higher rates for larger trees
<b>Evergreen Tree Crops</b>	Avocados, Bananas, Citrus, Olives	5-10L/ha or 1ml per sq mtr of plant area	Planting - drench or water in with 1:400 dilution. Apply from Bud initiation and at each growth flush 10 L/ha/month on mature trees, fertigated in every month that you irrigate
<b>Fruiting Annuals</b>	Capsicums, Cucurbits, Strawberries, Tomatoes, Pumpkins	5L/ha	Apply at seedling and transplanting and from 15cm growth. Every 10-14 days or 4 applications as required
<b>Leafy Vegetables</b>	Asparagus, Brassicas, Celery, Herbs, Lettuces	5L/ha	Planting out - 1:400 dilution (2.5 mL or 2.5L/1,000L water), for watering in of seedlings in a water wheel planter or similar device. Apply at seedling and transplanting and from 3-4 leaf. Every 10-14 days or 4 applications as required
<b>Root Vegetables</b>	Carrots, Onions, Potatoes, Sweet Potatoes	10L/ha	Apply from in furrow. 4-5 applications at critical growth stages
<b>Vine &amp; Berry Crops</b>	Blueberries, Raspberries, Wine and Table Grapes	5-10L/ha or 1ml/ sq mtr	Apply regularly during the growing season. As a minimum, use the product at times of plant stress or recovery such as Pre-Flowering, Fruit Filling & Post Harvest. Higher rates for larger canopies. <b>Do not apply directly over flowers where flowers are essential to crop productivity. Where bloom is a feature of fruit (e.g. blueberries &amp; table grapes) do not apply as a foliar spray once the bloom develops to avoid any damage</b>
<b>Broadacre</b>	Canola, Cotton, Pasture Crops (inc Lucerne), Sugar Cane, Turf	10L/ha or 1ml/ sq mtr	Apply at the start of the growing season and after each cut or grazing event in a rotational or strip grazing situation. Apply post emergence, 4-5 applications at critical growth stages. Seed coating (Cereals) Up to 4-10L/ tonne of seed. Do not exceed a total of 10L of total liquid per tonne. Sugar Cane - 5L/ha in furrow or billet dip, 2nd app 10L/ha at tillering. Ratoon crops 10L/ha at tillering

**Foliar Spraying:** Apply the product as soon as possible after mixing in the spray tank. Maintain agitation in spray tank if possible. **Seasol Plus Calcium** can be used as a foliar application or applied directly to the soil and can be applied with boom sprays, air blast sprays, drip systems, travelling irrigators, centre pivots and by aerial application.

**Fertigation:** Agitate the product regularly in the fertigation tank. For best results use the product on its own. **Seasol Plus Calcium** can be mixed with a number of agricultural chemicals, DO NOT COMBINE WITH CALCIUM NITRATE. See tank mix compatibility information at [www.seasol.com.au](http://www.seasol.com.au)

**Storage and Handling** Not to be kept for prolonged periods in hot conditions (>30°C) or in direct sunlight. Always use safe work practices for lifting and handling drums. Once diluted, the product should be applied within 24 hours. Agitate the product prior to using and re-agitate if the product is left standing for an extended period of time.

**Safety Directions:** Not to be taken. Keep out of reach of children. May irritate the skin and eyes. Avoid inhalation. Use in a well-ventilated space. Wash hands after use. Wash all edible plants before eating. If splashed, wash off with water. If swallowed or irritation persists, seek prompt medical advice. Additional information is listed in the Safety Data Sheet.

**Conditions of sale** This product must be used strictly in accordance to the directions. The efficacy of the product may be influenced by environmental conditions and application procedures and no warranty, express or implied is offered.



Scan QR code for further information. You can refer to our website, [www.seasol.com.au](http://www.seasol.com.au), or your local Seasol supplier for a range of Crop Bulletins and crop-specific Application Notes. You are also welcome to contact us during business hours on 1800 335 508 (toll-free within Aus).

**SEASOL INTERNATIONAL PTY LTD**  
1027 Mountain Highway, Bayswater, VIC, 3153

